



# DIGITAL ACCESS TO SCHOLARSHIP AT HARVARD

## Redefining Eosinophil Crystalloid Granules as a Potential New Functional Unit in Extracellular Inflammatory Events

The Harvard community has made this article openly available.  
[Please share](#) how this access benefits you. Your story matters.

Citation	Mahmudi-Azer, Salahaddin, Peter F. Weller, Ann M. Dvorak, Redwan Moqbel, and Peter D. Paré. 2010. Redefining eosinophil crystalloid granules as a potential new functional unit in extracellular inflammatory Events. Allergy, Asthma, and Clinical Immunology 6(Suppl. 2): P19.
Published Version	<a href="https://doi.org/10.1186/1710-1492-6-S2-P19">doi:10.1186/1710-1492-6-S2-P19</a>
Accessed	February 19, 2015 10:45:23 AM EST
Citable Link	<a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:10436092">http://nrs.harvard.edu/urn-3:HUL.InstRepos:10436092</a>
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA">http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA</a>

*(Article begins on next page)*



POSTER PRESENTATION

Open Access

# Redefining eosinophil crystalloid granules as a potential new functional unit in extracellular inflammatory Events

Salahaddin Mahmudi-Azer<sup>1\*</sup>, Peter F Weller<sup>2</sup>, Ann M Dvorak<sup>2</sup>, Redwan Moqbel<sup>3</sup>, Peter D Paré<sup>4</sup>

From Canadian Society of Allergy and Clinical Immunology Annual Scientific Meeting 2010  
Victoria, Canada. 3-6 November 2010

Eosinophils are major effector cells in allergic inflammatory response. They are known to synthesize, store, and release a wide range of pro-inflammatory mediators. Eosinophils contain different populations of mediator-storage organelles, including small secretory vesicles as well as crystalloid granules. In cytolysis, eosinophil cell membrane loses its integrity and crystalloid granules are released to extracellular space. Potential function of crystalloid granules in extracellular space as it relates to inflammatory events remains widely unknown. We hypothesized that eosinophil crystalloid granules are equipped to function independently in extracellular space. Our findings indicate that both DNA and RNA localize to human and rabbit eosinophil crystalloid granules and that RNA seems to be synthesized in intra-granular space further suggesting the presence of functional transcription machinery inside the granules. Furthermore, we show here that crystalloid granules express functional membrane receptors for a cytokine, IFN-gamma, as well as G protein-coupled membrane receptors for a chemokine, eotaxin. Our findings indicate that these receptors function by activating signal-transducing pathways within granules leading to mediator release from granules to extra-granular space in a cell free environment. Taken together our findings define a new potential role for eosinophil crystalloid granules as independent extracellular functional units in inflammatory events and may reveal a novel target in modulating the inflammatory events.

## Author details

<sup>1</sup>From Department of Medicine, University of Calgary, Alberta, Canada.

<sup>2</sup>Harvard Thorndike Laboratory and Charles A. Dana Research Institute,

\* Correspondence: sazer@shaw.ca

<sup>1</sup>From Department of Medicine, University of Calgary, Alberta, Canada

Full list of author information is available at the end of the article

Departments of Medicine and Pathology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215, USA. <sup>3</sup>Department of Immunology, University of Manitoba, Faculty of Medicine, Winnipeg, Manitoba, Canada. <sup>4</sup>The James Hogg iCAPTURE Centre, St. Paul's Hospital, Dept. of Medicine, University of British Columbia, Canada.

Published: 4 November 2010

doi:10.1186/1710-1492-6-S2-P19

**Cite this article as:** Mahmudi-Azer et al.: Redefining eosinophil crystalloid granules as a potential new functional unit in extracellular inflammatory Events. *Allergy, Asthma & Clinical Immunology* 2010 **6**(Suppl 2):P19.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

